




### Individual-days




### Aggressive acts to self




### Aggressive acts to others




### Individuals having alleged abuse, neglect or exploitation




### Body mass index




### Individuals with body mass index greater than 25 (overweight to obese)




### Body mass index change




### Weight change




### Waist circumference, males




### Combined pharmacotherapy




**Electroconvulsive treatment**  
(Each reported as zero for each of the months reported)



### Escape/Walkaway




### Falls




### Homicide

("Any homicide" and "any homicide attempt" both reported as zero incidents all months prior to September 2007)




### Incidents of individuals testing positive for illicit substance use


(Reported as zero incidents for each month )




### Medication variance (estimated)




### Medication variance (estimated)




### Medication variance (estimated)




### Medication variance (estimated)




### Medication variance (estimated)




**Any event involving a medication error which results in a major injury or exacerbation of a disease or disorder**




### Mortality




**Non-adherence to WRP for more than 20% of the interventions  
in seven consecutive days**



### Close observation




### Aspiration pneumonia




### Bowel Dysfunction


Individuals with occurrences of fecal impaction or bowel obstruction  
(Reported as zero for each of the months reported)




### Individuals with diagnosis of diabetes mellitus (estimated)




### Individuals with diagnosis of diabetes mellitus (estimated)




### Individuals with diagnosis of diabetes mellitus (estimated)




### Dysphagia




### Fractures



### Individuals with diagnosis of hepatitis C (estimated)




### Hospitalizations




### Individuals with diagnosis of MRSA


(Note: As of September 2007, count includes individuals whose infections have healed but who have carrier status.)




### Individuals with diagnosis of osteoporosis (estimated)




### Individuals with diagnosis of polydipsia (estimated)




### Seizure disorder




### Individuals receiving older anticonvulsants




### Skin integrity




### PRN medications




### Restraint



### Seclusion



### Stat medications



### Suicide

